




Affix patient label

<b>Arterial Line Care Plan</b>	
Insertion date/time:	
Insertion site:	
Inserted by:	

		PACU		Day 1		Day 2		Significant events/Comments
		Date:		Date:		Date:		
		Yes	No	Yes	No	Yes	No	
VIP Score	AM							
	PM							
	N							
Insertion site checked	AM							
	PM							
	N							
Patient apyrexial	AM							
	PM							
	N							
0.9% NCL flush bag and prescribed	AM							
	PM							
	N							
Pressure bag = 300mmHg	AM							
	PM							
	N							
Patient free of distal pain	AM							
	PM							
	N							
Distal sensation normal	AM							
	PM							
	N							
Normal capillary refill	AM							
	PM							
	N							
Normal colour	AM							
	PM							
	N							
Transparent dressing intact	AM							
	PM							
	N							
Non-injectable bung	AM							
	PM							
	N							
Patient board indicates art-line insitu	AM							
	PM							
	N							

<b>Line Removal</b>			
Date of removal:		No of days	
Reason for removal:		Tip sent:	
Name (print):		Signature	

**Enhanced Perioperative Protocol**



- 1) Fluid responsiveness is shown with a >10% increase in stroke volume after a passive leg raise or fluid bolus.
- 2) After four fluid boluses please take an ABG and seek a medical review
- 3) Ensure baseline ABG is taken prior to starting vasopressors. On the ward use the metaraminol protocol described in the Perioperative Manual and on the pharmacy monograph.

**Additional Plans/Deviations/Comments**



